

9803014
UTHEP-364
March, 1998

Exceptional Seiberg-Witten Geometry with Massive Fundamental Matters

Seiji Terashima and Sung-Kil Yang

Institute of Physics, University of Tsukuba, Ibaraki 305-0006, Japan

Abstract

We propose Seiberg-Witten geometry for $N = 2$ gauge theory with gauge group E_6 with massive N_f fundamental hypermultiplets. The relevant manifold is described as a fibration of the ALE space of E_6 type. It is observed that the fibering data over the base $\mathbf{CP}^1$ has an intricate dependence on hypermultiplet bare masses.

Recently several attempts have been made in extending the work of Seiberg and Witten on four-dimensional $N = 2$ supersymmetric gauge theory [1],[2] so as to include matter hypermultiplets in representations other than the fundamentals [3]-[6]. In our previous paper [3] we have refined the technique of $N = 1$ confining phase superpotentials toward the application to $N = 2$ or $N = 1$ gauge theories with A-D-E gauge groups with matter hypermultiplets in addition to an adjoint matter. The moduli space of $N = 1$ confining phase can also be studied in view of M -theory fivebrane, the result of which is in agreement with the field theory analysis [7],[8]. In [4] the fivebrane configurations are considered to obtain complex curves describing $N = 2$ $SU(N_c)$ gauge theory with matters in the symmetric or antisymmetric representation. In [5] the approach based on Type IIA string compactification on Calabi-Yau threefolds is used to find exact descriptions of $N = 2$ $SO(N_c)$ gauge theory with vectors and spinors.

In this paper we propose Seiberg-Witten geometry for $N = 2$ supersymmetric gauge theory with gauge group E_6 with massive N_f fundamental hypermultiplets. To this aim we employ the technique of $N = 1$ confining phase superpotentials [9]-[14]. The ALE space description of Seiberg-Witten geometry for $N = 2$ $SU(N_c)$ and $SO(2N_c)$ QCD is recovered in this approach [3]. We will show in the following that an extension of [3] enables us to obtain exceptional Seiberg-Witten geometry with fundamental hypermultiplets. The resulting manifold takes the form of a fibration of the ALE space of type E_6 .

Let us consider $N = 1$ E_6 gauge theory with N_f fundamental matters $Q^i, \tilde{Q}_j$ ($1 \leq i, j \leq N_f$) and an adjoint matter Φ . $Q^i, \tilde{Q}_j$ are in **27** and $\overline{\mathbf{27}}$, and Φ in **78** of E_6 . The coefficient of the one-loop beta function is given by $b = 24 - 6N_f$, and hence the theory is asymptotically free for $N_f = 0, 1, 2, 3$ and finite for $N_f = 4$. We take a tree-level superpotential

$$W = \sum_{k \in \mathcal{S}} g_k s_k(\Phi) + \text{Tr}_{N_f} \gamma_0 \tilde{Q}Q + \text{Tr}_{N_f} \gamma_1 \tilde{Q}\Phi Q, \quad (1)$$

where $\mathcal{S} = \{2, 5, 6, 8, 9, 12\}$ denotes the set of degrees of E_6 Casimirs $s_k(\Phi)$ and $g_k, (\gamma_a)_i^j$ ($1 \leq i, j \leq N_f$) are coupling constants. A basis for the E_6 Casimirs will be specified momentarily. When we put $(\gamma_0)_j^i = \sqrt{2}m_j^i$ with $[m, m^\dagger] = 0$, $(\gamma_1)_i^j = \sqrt{2}\delta_i^j$ and all $g_k = 0$, (1) is reduced to the superpotential in $N = 2$ supersymmetric Yang-Mills theory with massive N_f hypermultiplets.

We now look at the Coulomb phase with $Q = \tilde{Q} = 0$. Since Φ is restricted to take the values in the Cartan subalgebra we express the classical value of Φ in terms of a vector *

$$a = \sum_{i=1}^6 a_i \alpha_i \quad (2)$$

with α_i being the simple roots of E_6 . Then the classical vacuum is parametrized by

$$\Phi^{cl} = \text{diag}(a \cdot \lambda_1, a \cdot \lambda_2, \dots, a \cdot \lambda_{27}), \quad (3)$$

where λ_i are the weights for **27** of E_6 . For the notation of roots and weights we follow [15]. We define a basis for the E_6 Casimirs $u_k(\Phi)$ by

$$\begin{aligned} u_2 &= -\frac{1}{12} \chi_2, & u_5 &= -\frac{1}{60} \chi_5, & u_6 &= -\frac{1}{6} \chi_6 + \frac{1}{6 \cdot 12^2} \chi_2^3, \\ u_8 &= -\frac{1}{40} \chi_8 + \frac{1}{180} \chi_2 \chi_6 - \frac{1}{2 \cdot 12^4} \chi_2^4, & u_9 &= -\frac{1}{7 \cdot 6^2} \chi_9 + \frac{1}{20 \cdot 6^3} \chi_2^2 \chi_5, \\ u_{12} &= -\frac{1}{60} \chi_{12} + \frac{1}{5 \cdot 6^3} \chi_6^2 + \frac{13}{5 \cdot 12^3} \chi_2 \chi_5^2 \\ &\quad + \frac{5}{2 \cdot 12^3} \chi_2^2 \chi_8 - \frac{1}{3 \cdot 6^4} \chi_2^3 \chi_6 + \frac{29}{10 \cdot 12^6} \chi_2^6, \end{aligned} \quad (4)$$

where $\chi_n = \text{Tr } \Phi^n$. The standard basis $w_k(\Phi)$ are written in terms of u_k as follows

$$\begin{aligned} w_2 &= \frac{1}{2} u_2, & w_5 &= -\frac{1}{4} u_5, & w_6 &= \frac{1}{96} (u_6 - u_2^3), \\ w_8 &= \frac{1}{96} \left(u_8 + \frac{1}{4} u_2 u_6 - \frac{1}{8} u_2^4 \right), & w_9 &= -\frac{1}{48} \left(u_9 - \frac{1}{4} u_2^2 u_5 \right), \\ w_{12} &= \frac{1}{3456} \left(u_{12} + \frac{3}{32} u_6^2 - \frac{3}{4} u_2^2 u_8 - \frac{3}{16} u_2^3 u_6 + \frac{1}{16} u_2^6 \right). \end{aligned} \quad (5)$$

The basis $\{u_k\}$ and (5) were first introduced in [16].[†] In our superpotential (1) we then set

$$s_2 = w_2, \quad s_5 = w_5, \quad s_6 = w_6, \quad s_8 = w_8, \quad s_9 = w_9, \quad s_{12} = w_{12} - \frac{1}{4} w_6^2. \quad (6)$$

We will discuss later why this particular form is assumed.

The equations of motion are given by

$$\frac{\partial W(a)}{\partial a_i} = \sum_{k \in S} g_k \frac{\partial s_k(a)}{\partial a_i} = 0. \quad (7)$$

*Our notation is slightly different from [3]. Here we use a_i with lower index instead of a^i in [3].

[†]The Casimirs $u_1, u_2, u_3, u_4, u_5, u_6$ in [16] are denoted here as $u_2, u_5, u_6, u_8, u_9, u_{12}$, respectively.

Let us focus on the classical vacua with an unbroken $SU(2) \times U(1)^5$ gauge symmetry. Fix the $SU(2)$ direction by choosing the simple root α_1 , then we have the vacuum condition

$$a \cdot \alpha_1 = 2a_1 - a_2 = 0. \quad (8)$$

It follows from (7), (8) that

$$\begin{aligned} \frac{g_9}{g_{12}} &= \frac{D_{1,9}}{D_{1,12}} \\ &= -\frac{1}{8} \left(2a_1a_5a_4 - a_4a_3^2 + a_5^2a_4 + a_4^2a_3 - a_3a_6^2 + a_3^2a_6 \right. \\ &\quad \left. - 2a_1a_5^2 + 2a_1a_6^2 - 2a_4^2a_1 - a_5a_4^2 - 2a_1a_3a_6 + 2a_4a_3a_1 \right), \\ \frac{g_8}{g_{12}} &= \frac{D_{1,8}}{D_{1,12}} \\ &= -\frac{1}{48} \left(12a_1a_5^2a_4 - 6a_1^2a_5^2 - 6a_1^2a_6^2 - 4a_1^3a_3 + 4a_3^3a_1 + 2a_3^3a_4 + 2a_3^3a_6 - a_4^4 \right. \\ &\quad \left. - 3a_3^2a_6^2 - 3a_3^2a_4^2 - a_3^4 - a_6^4 - 12a_1a_5a_4^2 - 2a_5a_4a_6^2 + 8a_1a_3a_6^2 + 3a_1^4 \right. \\ &\quad \left. + 6a_1^2a_4a_3 - 8a_4a_3^2a_1 - 2a_5a_4a_3^2 - 2a_4^2a_3a_6 + 6a_1^2a_5a_4 - 2a_4a_3a_6^2 \right. \\ &\quad \left. + 2a_5a_4^2a_3 - 2a_5^2a_3a_6 + 2a_4a_3^2a_6 - a_5^4 - 2a_5^2a_4a_3 - 2a_1^2a_3^2 - 6a_1^2a_4^2 \right. \\ &\quad \left. + 2a_5^2a_6^2 + 2a_4^2a_6^2 + 2a_5^2a_3^2 + 6a_1^2a_3a_6 - 8a_1a_3^2a_6 + 8a_4^2a_3a_1 - 4a_5^2a_1a_3 \right. \\ &\quad \left. + 2a_5a_4a_3a_6 + 4a_5a_4a_1a_3 + 2a_3a_4^3 - 3a_4^2a_5^2 + 2a_4a_5^3 + 2a_4^3a_5 + 2a_6^3a_3 \right), \\ \frac{g_6}{g_{12}} &= \frac{D_{1,6}}{D_{1,12}} \\ &= \frac{1}{192} \left(4a_3^3a_1a_5^2 - 18a_4^3a_1^2a_5 + 13a_3^4a_1^2 - a_3^4a_5^2 - 7a_3^3a_6^3 + 9a_1^2a_6^4 + \dots \right), \end{aligned} \quad (9)$$

where $D_{1,k}$ is the cofactor for a $(1, k)$ element of the 6×6 matrix $[\partial s_i(a)/\partial a_j]$, $i \in \mathcal{S}$ and $j = 1, \dots, 6$ [3]. In (9) the explicit expression for g_6/g_{12} is too long to be presented here, and hence suppressed. Denoting $y_1 = g_9/g_{12}$, $y_2 = g_8/g_{12}$, $y_3 = g_6/g_{12}$, we find that the others are expressed in terms of y_1, y_2

$$\frac{g_2}{g_{12}} = \frac{D_{2,2}}{D_{2,12}} = y_1^2 y_2, \quad \frac{g_5}{g_{12}} = \frac{D_{2,5}}{D_{2,12}} = y_1 y_2. \quad (10)$$

This means that our superpotential specified with Casimirs (6) realizes the $SU(2) \times U(1)^5$ vacua only when the coupling constants are subject to the relation (10).

Notice that reading off degrees of y_1, y_2, y_3 from (9) gives $[y_1] = 3$, $[y_2] = 4$, $[y_3] = 6$. Thus, if we regard y_1, y_2, y_3 as variables to describe the E_6 singularity, (9) and (10) may

be identified as relevant monomials in versal deformations of the E_6 singularity. In fact we now point out an intimate relationship between classical solutions corresponding to the symmetry breaking $E_6 \supset SU(2) \times U(1)^5$ and the E_6 singularity. For this we examine the superpotential (1) at classical solutions

$$\begin{aligned} W_{cl} &= g_{12} \sum_{k \in S} \left(\frac{g_k}{g_{12}} \right) s_k^{cl}(a) \\ &= g_{12} \left(s_2^{cl} y_1^2 y_2 + s_5^{cl} y_1 y_2 + s_6^{cl} y_3 + s_8^{cl} y_2 + s_9^{cl} y_1 + s_{12}^{cl} \right). \end{aligned} \quad (11)$$

Evaluating the RHS with the use of (8)-(10) leads to

$$W_{cl} = -g_{12} \left(2y_1^2 y_3 + y_2^3 - y_3^2 \right). \quad (12)$$

It is also checked explicitly that

$$\begin{aligned} -4y_1 y_3 &= 2s_2^{cl} y_1 y_2 + s_5^{cl} y_2 + s_9^{cl}, \\ -3y_2^2 &= s_2^{cl} y_1^2 + s_5^{cl} y_1 + s_8^{cl}, \\ -2y_1^2 + 2y_3 &= s_6^{cl}. \end{aligned} \quad (13)$$

To illustrate the meaning of (11)-(13) let us recall the standard form of versal deformations of the E_6 singularity

$$W_{E_6}(x_1, x_2, x_3; w) = x_1^4 + x_2^3 + x_3^2 + w_2 x_1^2 x_2 + w_5 x_1 x_2 + w_6 x_1^2 + w_8 x_2 + w_9 x_1 + w_{12}, \quad (14)$$

where the deformation parameters w_k are related to the E_6 Casimirs via (5) [16]. Then what we have observed in (11)-(13) is that when we express w_k in terms of a_i as $w_k = w_k^{cl}(a)$ the equations

$$W_{E_6} = \frac{\partial W_{E_6}}{\partial x_1} = \frac{\partial W_{E_6}}{\partial x_2} = \frac{\partial W_{E_6}}{\partial x_3} = 0 \quad (15)$$

can be solved by [‡]

$$x_1 = y_1(a), \quad x_2 = y_2(a), \quad x_3 = i \left(y_3(a) - y_1(a)^2 - \frac{s_6^{cl}(a)}{2} \right) \quad (16)$$

under the condition (8). This observation plays a crucial role in our analysis.

[‡]We have observed a similar relation between the symmetry breaking solutions $SU(r+1)$ (or $SO(2r)$) $\supset SU(2) \times U(1)^{r-1}$ and the A_r (or D_r) singularity.

When applying the technique of confining phase superpotentials we usually take all coupling constants g_k as independent moduli parameters. To deal with $N = 1$ E_6 theory with fundamental matters, however, we find it appropriate to proceed as follows. First of all, motivated by the above observations for classical solutions, we keep three coupling constants $g'_6 = g_6/g_{12}$, $g'_8 = g_8/g_{12}$ and $g'_9 = g_9/g_{12}$ adjustable while the rest is fixed as $g'_2 = g'_8 g'^2_9$, $g'_5 = g'_8 g'_9$ with $g'_k = g_k/g_{12}$. Taking this parametrization it is seen that the equations of motion are satisfied by virtue of (10) in the $SU(2) \times U(1)^5$ vacua (8). Note here that originally there exist six classical moduli a_i among which one is fixed by (8) and three are converted to $g'_9 = y_1(a)$, $g'_8 = y_2(a)$ and $g'_6 = y_3(a)$, and hence we are left with two classical moduli which will be denoted as ξ_i . Without loss of generality one may choose $\xi_2 = s_2^{cl}(a)$ and $\xi_5 = s_5^{cl}(a)$.

We now evaluate the low-energy effective superpotential in the $SU(2) \times U(1)^5$ vacua. $U(1)$ photons decouple in the integrating-out process. The standard procedure yields the effective superpotential for low-energy $SU(2)$ theory [9],[3]

$$W_L = -g_{12} (2y_1^2 y_3 + y_2^3 - y_3^2) \pm 2\Lambda_{YM}^3, \quad (17)$$

where the second term takes account of $SU(2)$ gaugino condensation with Λ_{YM} being the dynamical scale for low-energy $SU(2)$ Yang-Mills theory. The low-energy scale Λ_{YM} is related to the high-energy scale Λ through the scale matching [3]

$$\begin{aligned} \Lambda_{YM}^6 &= g_{12}^2 A(a), \\ A(a) &\equiv \Lambda^{24-6N_f} \prod_{s=1}^6 \det_{N_f} (\gamma_0 + \gamma_1(a \cdot \lambda_s)), \end{aligned} \quad (18)$$

where λ_s are weights of **27** which branch to six $SU(2)$ doublets respectively under $E_6 \supset SU(2) \times U(1)^5$. Explicitly they are given in the Dynkin basis as

$$\begin{aligned} \lambda_1 &= (1, 0, 0, 0, 0, 0), & \lambda_2 &= (1, -1, 0, 0, 1, 0), \\ \lambda_3 &= (1, -1, 0, 1, -1, 0), & \lambda_4 &= (1, -1, 1, -1, 0, 0), \\ \lambda_5 &= (1, 0, -1, 0, 0, 1), & \lambda_6 &= (1, 0, 0, 0, 0, -1). \end{aligned} \quad (19)$$

Notice that $\sum_{s=1}^6 \lambda_s = 3\alpha_1$.

Let us first discuss the $N_f = 0$ case, i.e. E_6 pure Yang-Mills theory, for which $A(a)$ in (18) simply equals Λ^{24} . The vacuum expectation values are calculated from (17)

$$\begin{aligned}
\frac{\partial W_L}{\partial g_{12}} &= \langle \widetilde{W}(y_1, y_2, y_3; s) \rangle = -\left(2y_1^2 y_3 + y_2^3 - y_3^2\right) \pm 2\Lambda^{12}, \\
\frac{1}{g_{12}} \frac{\partial W_L}{\partial y_1} &= \left\langle \frac{\partial \widetilde{W}(y_1, y_2, y_3; s)}{\partial y_1} \right\rangle = -4y_1 y_3, \\
\frac{1}{g_{12}} \frac{\partial W_L}{\partial y_2} &= \left\langle \frac{\partial \widetilde{W}(y_1, y_2, y_3; s)}{\partial y_2} \right\rangle = -3y_2^2, \\
\frac{1}{g_{12}} \frac{\partial W_L}{\partial y_3} &= \left\langle \frac{\partial \widetilde{W}(y_1, y_2, y_3; s)}{\partial y_3} \right\rangle = -2y_1^2 + 2y_3,
\end{aligned} \tag{20}$$

where y_1, y_2, y_3 and g_{12} have been treated as independent parameters as discussed before and

$$\widetilde{W}(y_1, y_2, y_3; s) = s_2 y_1^2 y_2 + s_5 y_1 y_2 + s_6 y_3 + s_8 y_2 + s_9 y_1 + s_{12}. \tag{21}$$

Define a manifold by $\mathcal{W}_0 = 0$ with four coordinate variables $z, y_1, y_2, y_3 \in \mathbf{C}$ and

$$\mathcal{W}_0 \equiv z + \frac{\Lambda^{24}}{z} - \left(2y_1^2 y_3 + y_2^3 - y_3^2 + \widetilde{W}(y_1, y_2, y_3; s)\right) = 0. \tag{22}$$

It is easy to show that the expectation values (20) parametrize the singularities of the manifold where

$$\frac{\partial \mathcal{W}_0}{\partial z} = \frac{\partial \mathcal{W}_0}{\partial y_1} = \frac{\partial \mathcal{W}_0}{\partial y_2} = \frac{\partial \mathcal{W}_0}{\partial y_3} = 0. \tag{23}$$

Making a change of variables $y_1 = x_1, y_2 = x_2, y_3 = -ix_3 + x_1^2 + s_6/2$ in (22) we have

$$z + \frac{\Lambda^{24}}{z} - W_{E_6}(x_1, x_2, x_3; w) = 0. \tag{24}$$

Thus the ALE space description of $N = 2$ E_6 Yang-Mills theory [17],[16] is obtained from the $N = 1$ confining phase superpotential.

We next turn to considering the fundamental matters. In the $N = 2$ limit we have $A(a) = \Lambda^{24-6N_f} \cdot 8^{N_f} \prod_{i=1}^{N_f} f(a, m_i)$ with $f(a, m) = \prod_{s=1}^6 (m + a \cdot \lambda_s)$. After some algebra we find

$$f(a, m) = m^6 + 2\xi_2 m^4 - 8m^3 y_1 + \left(\xi_2^2 - 12y_2\right) m^2 + 4\xi_5 m - 4y_2 \xi_2 - 8y_3, \tag{25}$$

where we have used (6)-(9). Recall that, in viewing (17), we think of $(y_1, y_2, y_3, \xi_2, \xi_5, g_{12})$ as six independent parameters. Then the quantum expectation values are given by

$$\begin{aligned}\frac{\partial W_L}{\partial g_{12}} &= \langle \widetilde{W}(y_1, y_2, y_3; s) \rangle = -\left(2y_1^2 y_3 + y_2^3 - y_3^2\right) \pm 2\sqrt{A(y_1, y_2, y_3; \xi, m)}, \\ \frac{1}{g_{12}} \frac{\partial W_L}{\partial y_1} &= \left\langle \frac{\partial \widetilde{W}(y_1, y_2, y_3; s)}{\partial y_1} \right\rangle = -4y_1 y_3 \pm 2 \frac{\partial}{\partial y_1} \sqrt{A(y_1, y_2, y_3; \xi, m)}, \\ \frac{1}{g_{12}} \frac{\partial W_L}{\partial y_2} &= \left\langle \frac{\partial \widetilde{W}(y_1, y_2, y_3; s)}{\partial y_2} \right\rangle = -3y_2^2 \pm 2 \frac{\partial}{\partial y_2} \sqrt{A(y_1, y_2, y_3; \xi, m)}, \\ \frac{1}{g_{12}} \frac{\partial W_L}{\partial y_3} &= \left\langle \frac{\partial \widetilde{W}(y_1, y_2, y_3; s)}{\partial y_3} \right\rangle = -2y_1^2 + 2y_3 \pm 2 \frac{\partial}{\partial y_3} \sqrt{A(y_1, y_2, y_3; \xi, m)}.\end{aligned}\quad (26)$$

Similarly to the $N_f = 0$ case one can check that these expectation values satisfy the singularity condition for a manifold defined by

$$z + \frac{1}{z} A(y_1, y_2, y_3; \xi, m) - \left(2y_1^2 y_3 + y_2^3 - y_3^2 + \widetilde{W}(y_1, y_2, y_3; s)\right) = 0. \quad (27)$$

Note that s_k in $\widetilde{W}$ are quantum moduli parameters. What about ξ_2, ξ_5 in the one-instanton factor A ? Classically we have $\xi_i = s_i^{cl}$ as was seen before. The issue is thus whether the classical relations $\xi_i = s_i^{cl}$ receive any quantum corrections at the singularities. If there appear no quantum corrections, ξ_i in A can be replaced by quantum moduli parameters s_i . Let us simply assume here that $\xi_i = s_i^{cl} = \langle s_i \rangle$ for $i = 2, 5$ in the $N = 1$ $SU(2) \times U(1)^5$ vacua. This assumption seems quite plausible as long as we have inspected possible forms of quantum corrections due to gaugino condensates.

Now we find that Seiberg-Witten geometry of $N = 2$ supersymmetric QCD with gauge group E_6 is described by

$$z + \frac{1}{z} A(x_1, x_2, x_3; w, m) - W_{E_6}(x_1, x_2, x_3; w) = 0, \quad (28)$$

where a change of variables from y_i to x_i as in (24) has been made in (27) and

$$\begin{aligned}& A(x_1, x_2, x_3; w, m) \\ &= \Lambda^{24-6N_f} \cdot 8^{N_f} \prod_{i=1}^{N_f} \left(m_i^6 + 2w_2 m_i^4 - 8m_i^3 x_1 + (w_2^2 - 12x_2) m_i^2 \right. \\ &\quad \left. + 4w_5 m_i - 4w_2 y_2 - 8(x_1^2 - ix_3 + w_6/2) \right).\end{aligned}\quad (29)$$

The manifold takes the form of ALE space of type E_6 fibered over the base $\mathbf{CP}^1$. Note an intricate dependence of the fibering data over $\mathbf{CP}^1$ on the hypermultiplet masses. This is

in contrast with the ALE space description of $N = 2$ $SU(N_c)$ and $SO(2N_c)$ gauge theories with fundamental matters. In (28), letting $m_i \rightarrow \infty$ while keeping $\Lambda^{24-6N_f} \prod_{i=1}^{N_f} m_i^6 \equiv \Lambda_0^{24}$ finite we recover the pure Yang-Mills result (24).

As a non-trivial check of our proposal (28) let us examine the semi-classical singularities. In the semi-classical limit $\Lambda \rightarrow 0$ the discriminant Δ for (28) is expected to take the form $\Delta \propto \Delta_G \Delta_M$ where Δ_G is a piece arising from the classical singularities associated with the gauge symmetry enhancement and Δ_M represents the semi-classical singularities at which squarks become massless. When the N_f matter hypermultiplets belong to the representation $\mathcal{R}$ of the gauge group G we have

$$\Delta_M = \prod_{i=1}^{N_f} \det_{d \times d}(m_i \mathbf{1} - \Phi^{cl}) = \prod_{i=1}^{N_f} P_G^{\mathcal{R}}(m_i; u), \quad (30)$$

where $d = \dim \mathcal{R}$, m_i are the masses, Φ^{cl} denotes the classical Higgs expectation values and $P_G^{\mathcal{R}}(x; u)$ is the characteristic polynomial for $\mathcal{R}$ with u_i being Casimirs constructed from Φ^{cl} .

For simplicity, let us consider the case in which all the quarks have equal bare masses. Then we can change a variable x_3 to $\tilde{x}_3$ so that $A = A(\tilde{x}_3; w, m)$ is independent of x_1 and x_2 . Eliminating x_1 and x_2 from (28) by the use of

$$\frac{\partial W_{E_6}}{\partial x_1} = \frac{\partial W_{E_6}}{\partial x_2} = 0, \quad (31)$$

we obtain a curve which is singular at the discriminant locus of (28). The curve is implicitly defined through

$$\overline{W}_{E_6} \left(\tilde{x}_3; w_i - \delta_{i,12} \left(z + \frac{A(\tilde{x}_3; w, m)}{z} \right) \right) = 0, \quad (32)$$

where $\overline{W}_{E_6}(\tilde{x}_3; w_i) = W_{E_6}(x_1(\tilde{x}_3, w_i), x_2(\tilde{x}_3, w_i), \tilde{x}_3; w_i)$ and $x_1(\tilde{x}_3, w_i)$, $x_2(\tilde{x}_3, w_i)$ are solutions of (31). Now the values of $\tilde{x}_3$ and z at singularities of this curve can be expanded in powers of $\Lambda^{\frac{24-6N_f}{2}}$. Then it is more or less clear that the classical singularities corresponding to massless gauge bosons are produced. Furthermore, if we denote as $R(\overline{W}, A)$ the resultant of $\overline{W}_{E_6}(\tilde{x}_3; w_i)$ and $A(\tilde{x}_3; w, m)$, then $R(\overline{W}, A) = 0$ yields another singularity condition of the curve in the limit $\Lambda \rightarrow 0$. We expect that $R(\overline{W}, A) = 0$ corresponds to the semi-classical massless squark singularities as is observed in the case

of $N = 2$ $SU(N_c)$ QCD [18],[14]. Indeed, we have checked this by explicitly computing $R(\overline{W}, A)$ at sufficiently many points in the moduli space. For instance, taking $w_2 = 2, w_5 = 5, w_6 = 7, w_8 = 9, w_9 = 11$ and $w_{12} = 13$ in the $N_f = 1$ case, we get

$$\begin{aligned}
& R(\overline{W}, A) \\
&= m^2 \left(3m^{10} + 12m^8 + \dots \right) \left(26973m^{27} + 258552m^{25} + \dots \right)^3 \\
&\quad \left(m^{27} + 24m^{25} + 240m^{23} + 240m^{22} + 2016m^{21} + 3360m^{20} + 16416m^{19} \right. \\
&\quad + 34944m^{18} + 88080m^{17} + 216576m^{16} + 448864m^{15} + 607488m^{14} \\
&\quad + 2198272m^{13} - 296000m^{12} + 4177792m^{11} - 3407104m^{10} + 7796224m^9 \\
&\quad + 10664448m^8 - 31708160m^7 + 41183232m^6 - 21889792m^5 + 15575040m^4 \\
&\quad \left. - 17125120m^3 - 38456320m^2 - 3461120m + 9798656 \right), \tag{33}
\end{aligned}$$

while the E_6 characteristic polynomial for **27** is given by

$$\begin{aligned}
& P_{E_6}^{\mathbf{27}}(x; u) \\
&= x^{27} + 12w_2x^{25} + 60w_2^2x^{23} + 48w_5x^{22} + (96w_6 + 168w_2^3)x^{21} + 336w_2w_5x^{20} \\
&\quad + (528w_2w_6 + 294w_2^4 + 480w_8)x^{19} + (1344w_9 + 1008w_2^2w_5)x^{18} + \dots \tag{34}
\end{aligned}$$

We now find a remarkable result that the last factor of (33) precisely coincides with $P_{E_6}^{\mathbf{27}}(m; u)$! Hence the manifold described by (28) correctly produces all the semi-classical singularities in the moduli space of $N = 2$ supersymmetric E_6 QCD.

If we choose another form of the superpotential (1), say, the superpotential with $s_i = w_i$ for $i \in \mathcal{S}$ instead of (6) we are unable to obtain Δ_M in (30). As long as we have checked the choice made in (6) is judicious in order to pass the semi-classical test. At present, we have no definite recipe to fix the tree-level superpotential which produces the correct semi-classical singularities, though it is possible to proceed by trial and error. In fact we can find Seiberg-Witten geometry for $N = 2$ $SO(2N_c)$ gauge theory with spinor matters and $N = 2$ $SU(N_c)$ gauge theory with antisymmetric matters [19].

In our result (28) it may be worth mentioning that the gaussian variable x_3 of the E_6 singularity appears in the fibering term. An analogous structure is observed for $N = 2$ $SO(10)$ gauge theory with spinors and vectors in [5]. Their result reads

$$z + \frac{\Lambda^{2b}}{z} B(x_1, x_3; v) - W_{D_5}(x_1, x_2, x_3; v) = 0, \tag{35}$$

where $b = 8 - (6 - n) - 2(4 - n) = 3n - 6$ is the coefficient of the one-loop beta function and

$$\begin{aligned} W_{D_5}(x_1, x_2, x_3; v) &= x_1^4 + x_1 x_2^2 - x_3^2 + v_2 x_1^3 + v_4 x_1^2 + v_6 x_1 + v_8 + v_5 x_2, \\ B(x_1, x_3; v) &= x_1^{6-n} \left(x_3 - \frac{1}{8} (4v_4 - v_2^2 + 4v_2 x_1 + 8x_1^2) \right)^{4-n} \end{aligned} \quad (36)$$

for $SO(10)$ with massless $(6 - n)$ fundamentals and $(4 - n)$ spinors. Here v_i stand for the $SO(10)$ Casimirs. It is tempting to suspect that the above E_6 and $SO(10)$ results are related through the Higgs mechanism under the symmetry breaking $E_6 \supset SO(10) \times U(1)$.

There is no difficulty in using our method to find Seiberg-Witten geometry in the form of ALE fibrations for $N = 2$ QCD with E_7 gauge group although more computer powers are obviously required. Finally, to compare the present results, it is desired to work out exceptional Seiberg-Witten geometry with fundamental matters in the framework of Calabi-Yau geometric engineering [20]-[22].

The work of S.T. is supported by JSPS Research Fellowship for Young Scientists. The work of S.K.Y. was supported in part by the Grant-in-Aid for Scientific Research on Priority Area “Physics of CP violation”, the Ministry of Education, Science and Culture, Japan.

References

- [1] N. Seiberg and E. Witten, Nucl. Phys. **B426** (1994) 19, hep-th/9407087; Nucl. Phys. **B431** (1994) 484, hep-th/9408099
- [2] For recent reviews, see, W. Lerche, *Introduction to Seiberg-Witten Geometry and its Stringy Origin*, Nucl. Phys. Proc. Suppl. **55B** (1997) 83, Fortsch. Phys. **45** (1997) 293, hep-th/9611190; A. Klemm, *On the Geometry behind $N=2$ Supersymmetric Effective Actions in Four Dimensions*, hep-th/9705131
- [3] S. Terashima and S.-K. Yang, *ADE Confining Phase Superpotentials*, hep-th/9706076, to appear in Nucl. Phys. **B**
- [4] K. Landsteiner and E. Lopez, *New Curves from Branes*, hep-th/9708118
- [5] M. Aganagic and M. Gremm, *Exact Solutions for Some $N=2$ supersymmetric $SO(N)$ gauge theories with vectors and spinors*, hep-th/9712011
- [6] H. Oda, S. Tomizawa, N. Sakai and T. Sakai, *M theory and the “Integrating in” Method with an Antisymmetric Tensor*, hep-th/9802004
- [7] J. de Boer and Y. Oz, *Monopole Condensation and Confining Phase of $N = 1$ Gauge Theories via M Theory Five-Brane*, hep-th/9708044
- [8] S. Terashima, *Supersymmetric Gauge Theories with Classical Groups via M Theory Five-Brane*, hep-th/9712172
- [9] S. Elitzur, A. Forge, A. Giveon, K. Intriligator and E. Rabinovici, Phys. Lett. **B379** (1996) 121, hep-th/9603051
- [10] S. Terashima and S.-K. Yang, Phys. Lett. **B391** (1997) 107, hep-th/9607151
- [11] K. Landsteiner, J.M. Pierre and S.B. Giddings, Phys. Rev. **D55** (1997) 2367, hep-th/9609059
- [12] T. Kitao, Phys. Lett. **B402** (1997) 290, hep-th/9611097

- [13] T. Kitao, S. Terashima and S.-K. Yang, Phys. Lett. **B399** (1997) 75, hep-th/9701009
- [14] A. Giveon, O. Pelc and E. Rabinovici, Nucl. Phys. **B499** (1997) 100, hep-th/9701045
- [15] R. Slansky, Phys. Reports **79** (1981) 1
- [16] W. Lerche and N.P. Warner, *Exceptional SW Geometry from ALE Fibrations*, hep-th/9608183
- [17] A. Klemm, W. Lerche, P. Mayr, C. Vafa and N. Warner, Nucl. Phys. **B477** (1996) 746, hep-th/9604034
- [18] A. Hanany and Y. Oz, Nucl. Phys. **B452** (1995) 283, hep-th/9505075
- [19] S. Terashima and S.-K. Yang, to appear
- [20] S. Katz, A. Klemm and C. Vafa, Nucl. Phys. **B497** (1997) 173, hep-th/9609239
- [21] J.H. Brodie, Nucl. Phys. **B506** (1997) 183, hep-th/9705068
- [22] S. Katz, P. Mayr and C. Vafa, *Mirror Symmetry and Exact Solutions of 4D N=2 Gauge Theories-I*, hep-th/9706110